

HOME GARDEN TREATMENT RECOMMENDATIONS				
Crops	Pests Controlled	Maximum Number of Applications per Season	Minimum Days to Wait Before Reapplying	Minimum Days to Wait From Last Application to Harvest
cole crops (Brassica vegetables), including, but not limited to: broccoli, Chinese broccoli, broccoli raab, brussels sprouts, cabbage, Chinese cabbage (bok choy), Chinese cabbage (napa), cauliflower, cavalo, collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, Chinese mustard cabbage (gai choy) and rape greens	cabbage looper diamondback moth imported cabbage worm leafminers worms	6	4	1
cucurbits, including, but not limited to: cucumber, summer and winter squash, muskmelons (cantaloupe, honeydew, etc.), pumpkin, edible gourds and watermelon	leafminers thrips worms (caterpillars)	6	5	all except cucumber: 3 cucumber: 1
fruiting vegetables, including, but not limited to: eggplant, ground cherry, okra, pepino, pepper, tomatillo, and tomato	Colorado potato beetle leafminers thrips worms (caterpillars)	6	4	1

HOME GARDEN TREATMENT RECOMMENDATIONS				
Crops	Pests Controlled	Maximum Number of Applications per Season	Minimum Days to Wait Before Reapplying	Minimum Days to Wait From Last Application to Harvest
bushberries and caneberries, including, but not limited to: blueberry, blackberry, raspberry, orangeberry, currant, gooseberry, huckleberry, elderberry, juniperberry, lingonberry, and salal	armyworms fireworms fruitworms leafrollers looper thrips fruitfly (suppression)	6	6	3
citrus trees, including, but not limited to: oranges, grapefruit, lemons, limes, and tangerines (to prevent fruit scarring from thrips, treat when fruit is marble size)	katydids leafminers thrips worms (caterpillars)	6	6	1

Unit of Measure [†]	AMOUNT OF NATURAL-GUARD [®] SPINOSAD TO USE PER PINT, QUART OR GALLON OF SPRAY		
	Per Pint (16 Fl. Oz.) of Spray	Per Quart (32 Fl. Oz.) of Spray	Per Gallon (128 Fl. Oz.) of Spray
Fluid Ounces (Fl. Oz.)	0.25 Fl. Oz.	0.5 Fl. Oz.	2.0 Fl. Oz.
Milliliters (ml.)	7.5 ml.	15 ml.	60 ml.
Tablespoons (Tbs.)	1/2 Tbs.	1 Tbs.	4 Tbs.
Teaspoons (tsp.)	1 1/2 Tsp.	3 Tsp.	12 Tsp.

[†]Conversion factors: 1 Fl. Oz. = 30 ml. = 2 tablespoons (Tbs.) = 6 teaspoons (tsp.) (1 teaspoon = 1/2 tablespoon)

WHEN TO APPLY

Apply when listed pests are present. Repeat applications may be made as indicated in the Home Garden Treatment Recommendations table. See your state extension service recommendations for treatment guidelines in your area.

HOME GARDEN TREATMENT RECOMMENDATIONS				
Crops	Pests Controlled	Maximum Number of Applications per Season	Minimum Days to Wait Before Reapplying	Minimum Days to Wait From Last Application to Harvest
apple and other pome fruits including, but not limited to: pears, crabapples, mayhaw and quince	codling moth leafminers leafrollers Oriental fruit moth tufted apple budmoth	6	10	7
asparagus (post-harvest to protect ferns)	asparagus beetles	4	7	60

Distributed By:
VOLUNTARY PURCHASING GROUPS INC.
 P. O. BOX 460
BONHAM, TEXAS 75418
 EPA Reg. No. 62719-314-7401
 EPA Est. No. 7401-TX-1



WARRANTY DISCLAIMER

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS NOR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by Buyer.

LIMITATION OF REMEDIES

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

LAWN AND ORNAMENTAL RECOMMENDATIONS		
Use Site	Pests Controlled	Comments
outdoor ornamentals (herbaceous and woody plants)	gall midges leaf feeding beetles leafminers sawfly larvae spider mites ¹ thrips worms, including loopers, webworms, bagworms, gypsy moth, and tent caterpillars	Mix the amount of Natural-Guard[®] SPINOSAD recommended per pint, quart or gallon of spray and uniformly spray foliage to point of runoff. Uniform coverage of upper and lower leaf surfaces is essential for effective insect control.
lawns, ornamentals, home gardens (see listed crops in previous table) and other outdoor areas	fire ants	Individual Fire Ant Mounds: Mix the amount of Natural-Guard[®] SPINOSAD recommended per gallon of spray and use a sprinkler can or similar device to apply the diluted spray as a mound drench, using 1 to 2 gallons per mound. Do not use pressurized sprays. Apply about 10% of the spray volume around the perimeter of the mound and the remainder directly to the mound. Do not disturb fire ant mound prior to application. If possible, apply following recent rainfall. For best results apply when weather is cool, 65 to 85°F, or in early morning or late evening.

¹Apply when spider mites are first observed prior to webbing and before mite populations have become severe. Reapply after 7 to 10 days (3 to 5 days in greenhouses and structures that can be closed or open) to contact newly hatched nymphs and repeat until infestation is managed. **Uniform coverage of both upper and lower leaf surfaces is critical.** Addition of a nonionic spray adjuvant at 0.1% v/v has been shown to enhance control of spider mites (follow surfactant manufacturer's label recommendations).

TERMS AND CONDITIONS OF USE

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

The registrant acknowledges that this product may be misbranded if one or more of its ingredients, active and/or inert, is reclassified as non-organic and/or removed from the United States Department of Agriculture (USDA) National List of approved substances for use in organic production and handling. If one or more of the ingredients in this pesticide product are so reclassified and/or removed (delisted) from the National List, the registrant will take appropriate remedial measures with respect to the labeling for this product.

HOME GARDEN TREATMENT RECOMMENDATIONS				
Crops	Pests Controlled	Maximum Number of Applications per Season	Minimum Days to Wait Before Reapplying	Minimum Days to Wait From Last Application to Harvest
tuberous and corn vegetables, including, but not limited to: potatoes, sweet potatoes, yams, Jerusalem artichoke, Chinese artichoke, Cassava, chayote root, ginger, and turmeric	Colorado potato beetle, corn borers, leaf miners, thrips	6	7	7
all crops listed in this table		fire ants	See application directions in following table	

LAWN AND ORNAMENTAL RECOMMENDATIONS		
Use Site	Pests Controlled	Comments
lawns	worms, including small (< 1/4") armyworm, and sod webworm cat fleas (suppression only)	Mix the amount of Natural-Guard[®] SPINOSAD recommended per gallon of spray and uniformly apply a minimum of 3 gallons of spray per 1000 sq ft of area. Delay watering or mowing for 12 to 24 hours after application. Armyworms: For best results, apply in early morning or late afternoon. Cat fleas: Apply early or late in the day since control requires contact with dilute spray before drying. Thorough coverage is necessary. Reapply in 7 to 14 days to control adults that have emerged from pupae present at time of initial treatment.

HOME GARDEN TREATMENT RECOMMENDATIONS				
Crops	Pests Controlled	Maximum Number of Applications per Season	Minimum Days to Wait Before Reapplying	Minimum Days to Wait From Last Application to Harvest
succulent beans and peas , including, but not limited to: lima bean, snap bean, blackeyed pea, garden pea, garbanzo bean, and snow pea	borers leafminers thrips worms (caterpillars)	6	5	3
sweet corn and popcorn (for earworms, treat silk frequently as it grows)	corn borers worms (caterpillars), including earworm	6	3	1
tree nuts , including, but not limited to: pecans, almonds, cashew, chestnut, macadamia, pistachio, and walnuts	leafrollers navel orangeworms peach twig borer shuckworms webworms husk fly (suppression)	6	7	14
tropical tree fruits : atemoya, avocado, sapote, cherimoya, custard apple, guava, longan, lychee, mango, papaya, passionfruit, rambutan, sapodilla, star apple, starfruit, and wax jambu	suppression of worms or caterpillars, thrips, and katydids	4	7	1

HOME GARDEN TREATMENT RECOMMENDATIONS				
Crops	Pests Controlled	Maximum Number of Applications per Season	Minimum Days to Wait Before Reapplying	Minimum Days to Wait From Last Application to Harvest
grape	leafrollers thrips worms (berry moth)	6	5	7
leafy and dried herbs	leafminers thrips worms (caterpillars)	5	7	1
leafy vegetables , including, but not limited to: head and leaf lettuce, celery, arugula, chervil, edible chrysanthemum, corn salad, cress, dandelion, dock, endive, fennel, parsley, garden purslane, radicchio, rhubarb, spinach, and Swiss chard	leafminers worms (caterpillars)	6	4	1
stone fruits , including, but not limited to: peaches, plums, cherries, nectarines, prunes, and apricots	borers fruit flies leafminers leafrollers thrips worms (caterpillars)	6	7	cherry, plum and prune, 7 peach, nectarine and apricot, 14
strawberry	armyworms leafrollers thrips	6	5	1

Natural Guard®

GARDENING NATURE'S WAY



READY TO SPRAY

ACTIVE INGREDIENT:
Spinosad (a mixture of spinosyn
A and spinosyn D) 0.5%
INERT INGREDIENTS 99.5%
TOTAL 100.0%
Contains 0.04 lb of active ingredient per gallon.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label. If terms are unacceptable, return at once unopened. In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Shake Well Before Use –
Avoid Freezing

KEEP OUT OF REACH
OF CHILDREN

CAUTION

See Back Panel For
Additional Precautionary
Statements And Directions
For Use Including
Storage And Disposal.

SPINOSAD

LANDSCAPE AND GARDEN INSECTICIDE

FOR RESIDENTIAL USE IN HOME GARDENS, LAWN AND ORNAMENTALS
For Control Of Foliage Feeding Worms (Caterpillars), Thrips And Other Listed Pests In:

- Fruiting Vegetables, Cucurbits, Cole Crops, Leafy And Tuberous Vegetables, Stone Fruits, Citrus Trees, And Lawns

NET CONTENTS 32 FL. OZS.

Natural-Guard® SPINOSAD

25M-1-74

PRECAUTIONARY STATEMENTS ENVIRONMENTAL HAZARDS

This product is toxic to bees exposed to treatment for 3 hours following treatment. Do not apply this pesticide to blooming, pollen-shedding or nectar-producing parts of plants if bees may forage on the plants during this time period. This product is toxic to aquatic invertebrates. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not put product or rinse water down drain.

PHYSICAL OR CHEMICAL HAZARDS

COMBUSTIBLE. Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

For residential use in home gardens, lawn and ornamentals. Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container only in secure or locked storage area.

PESTICIDE DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800-992-3994 for disposal instructions. Never place unused product down any indoor or outdoor drain.

Natural-Guard® SPINOSAD is intended for control of worms (caterpillars) and other listed insects. Natural-Guard® SPINOSAD does not significantly impact predatory beneficial insects, predatory mites, and spiders while controlling target pests. Susceptible insect pests may be observed on plants up to several hours after treatment, but will have ceased active feeding before being killed. Natural-Guard® SPINOSAD is classified as an organic substance by the USDA National Organic Standards Board.

MIXING AND APPLICATION (SHAKE WELL BEFORE USE)

Natural-Guard® SPINOSAD may be applied with trigger sprayer, hand-held, backpack, or hose-end sprayers. Use a hose-end sprayer that can be adjusted to provide a dilution ratio of about 2 fl. oz. of Natural-Guard® SPINOSAD (4 Tbs.) per gallon of spray. See Hose-End Sprayer Directions below if this product is packaged in a hose-end sprayer. Add the required amount of Natural-Guard® SPINOSAD to the recommended amount of water, mix thoroughly, and apply uniformly to both upper and lower surfaces of plant foliage. It is recommended to mix only as much spray as needed for a single treatment. In vegetable gardens it is recommended to use not more than 3 gallons of spray for 1000 sq. ft. of area. Do not use kitchen utensils for measuring. Keep measuring utensils with product and away from children.

▼ PEEL FROM CORNER OF BOOKLET