

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a cool place. Activity may be impaired by storage at temperatures above 90°F.

PESTICIDE AND CONTAINER DISPOSAL:

If Empty: do not reuse this container. Place in trash or offer for recycling if available. If Partly Filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

Do not reuse container. Wrap container and put in trash.

CAUTION: Harmful if swallowed or inhaled. Avoid breathing vapors or spray mist. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

FIRST AID

If inhaled: Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center of doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If swallowed: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person. Have the product container of label with you when calling a poison control center or doctor, or going for treatment. Hot Line Number: 1-800-255-3924

ENVIRONMENTAL HAZARDS

Do not contaminate water when disposing of equipment washwaters or rinsate. Do not apply directly to water.

Seller warrants that the material contained herein conforms to the description on the label and is reasonably fit for the purposes referred to in the directions for use. Timing and method of application, weather, watering practices, nature of soil, the insect problem, condition of the crop, incompatibility with other chemical not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herein. NO OTHER EXPRESS OR IMPLIED WARRANTY OF THE FITNESS OR MERCHANTABILITY IS MADE.



Thuricide® Concentrate

- Controls: Gypsy Moth, Cankerworm, Tent Caterpillar, Cabbage Looper, Imported Cabbage Worm And Tomato Hornworm.
- Mix 4 Teaspoons Per Gallon Of Water For Shade Trees And Ornamentals.

ACTIVE INGREDIENT:

Bacillus thuringiensis subspecies *kurstaki*, Spores and Lepidopteran active toxins (at least 6 million viable spores per milligram*) 15%

OTHER INGREDIENTS:** 85%

TOTAL 100%

The percent active ingredient does not indicate product performance and potency measurements are not federally standardized.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**
See Side Panel For Additional Precautionary Statements

NET CONTENTS ONE PINT (16 Fl.Ozs.)

Distributed By:
VOLUNTARY PURCHASING GROUPS, INC.
P. O. BOX 460
BONHAM, TEXAS 75418
Visit Us At: www.v-p-g.com
EPA Reg. No. 70051-65-7401
EPA Est. No. 7401-TX-1



Thuricide® is a Registered Trademark of Certis USA, L.L.C.
20M-10-96K

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

HOW TO APPLY

Always shake or stir American Brand Concentrate thoroughly before use. Partially fill sprayer with water before adding the application rate amount of product, then mix in product thoroughly and add remaining amount of water. Spray leaf surfaces thoroughly, top and bottom, for complete control. Agitate regularly while spraying. Use all of spray mixture within 24 hours.

SHADE TREES AND ORNAMENTALS

Mix 2 fl. ozs. per 3 gallons of water (4 tsp. per gallon) for a handsprayer or proportional amount for a hose-end sprayer following Manufacturer's Directions. Apply at first sign of infestation and repeat at weekly intervals when needed to maintain control. Apply to thoroughly cover all foliage surfaces of Shade Trees and Ornamentals for control of the following leaf-feeding Worms:

Bagworm	Elm Spanworm	GypsyMoth(N.E.)*
Fall Cankerworm	Tent Caterpillar	Spring Cankerworm
Fall Webworm		

*Apply when buds swell, but before leaves are fully open. If eggs hatch over a long period of time or if reinfestation occurs, spray about 14 days after first application.

FRUITS AND VEGETABLES

Mix 1½ fl. ozs. per 3 gallons of water (1 Tbsp./Gallon) for a hand sprayer and apply to all plant foliage to cover an area of 1,000 square feet. Apply at first sign of infestation and repeat at weekly intervals when needed to maintain control of the following Insects: **CABBAGE LOOPER** on Broccoli, Cauliflower, Collards, Kale, Mustard Greens, Turnip Greens, Cabbage, Celery, Lettuce, Melons and Tomatoes. **IMPORTED CABBAGE WORMS** on Broccoli, Cabbage, Cauliflower, Collards, Kale, Mustard Greens and Turnip Greens. **TOMATO HORNWORM** on Tomatoes.