

Weed-Out Plus Lawn Fertilizer

Weed-Out Plus Lawn Fertilizer



ferti-lome®



Weed-Out PLUS LAWN FERTILIZER

CONTAINS *TRIMEC® HERBICIDE

- Controls Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge, And Many Other Broadleaf Weeds Which Are Listed On This Label In St. Augustinegrass, Centipedegrass, Bahiagrass, Bermudagrass, Zoysiagrass, Bluegrass, Perennial Ryegrass, Fescue and Bentgrass Lawns.



Covers
Up To
10,000
Sq. Ft.

ACTIVE INGREDIENTS:

Dimethylamine salt of (+)-2-(2-methyl-4-chlorophenoxy) propionic acid 0.29%

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid 0.175%

Dimethylamine salt of dicamba (3,6-dichloro-o-aminic acid) 0.07%

INERT INGREDIENTS:

99.47%

TOTAL: 100.00%

THIS PRODUCT CONTAINS:

4.80 lbs. or 0.24% (+)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton.

2.80 lbs. or 0.14% 2,4-dichlorophenoxyacetic acid equivalent per ton.

1.20 lbs. or 0.06% 3,6-dichloro-o-aminic acid equivalent per ton.

Isomer Specific by AOMC Methodology

Contract Manufacturing from Meuronon-p

TRIMEC® is a registered trademark of PB/GORDON CORPORATION.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for First Aid,
Precautionary Statements,
and Directions for Use.

NET WEIGHT 40 LBS. (18.14 KG)

10923

Weed-Out Plus Lawn Fertilizer



Weed-Out Plus Lawn Fertilizer



Weed-Out Plus Lawn Fertilizer

STOP! READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

FIRST AID

If In Eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.
If On Skin Or Clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a Poison Control Center or doctor for treatment advice.
If Swallowed:	<ul style="list-style-type: none"> Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment. You may also call toll free 1-877-800-5556 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from target area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried or dust has settled.

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

CONTAINER DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partially filled: Call your local solid waste agency or toll free number (1-800-CLEANUP) for disposal instructions. Never place unused product down any indoor or outdoor drain.

USE PRECAUTIONS

Do not use this product where desirable clovers are present. Do not apply in areas underlaid by roots of desirable trees or shrubs. Do not exceed specified dosages for any area. Do not apply to newly seeded lawns. During application, close spreader openings when turning. Do not contaminate domestic or irrigation waters.

INSTRUCTIONS

ferti-lome® WEED-OUT® PLUS LAWN FERTILIZER controls a wide range of Lawn Weeds, including Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and many others commonly found in Home Lawns. In addition, a 25-0-4 fertilizer provides a supply of plant food. Possible retreatment may be necessary if (a) Lawn is heavily overrun by Weeds or (b) if adverse low moisture conditions prevail and Weeds are in a state of poor growth. A second application may be made in approximately 30 days. Do not use on Dichondra. Do not over apply. Under conditions of adequate moisture, high temperatures and high humidity, over-application may cause minor transitory discoloration on St. Augustinegrass. Do not use this product on "Floratam" St. Augustinegrass.

The maximum application rate to turf is 0.3 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget areas.
4. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

HOW MUCH TO APPLY

For Kentucky Bluegrass, Fescues, Perennial Ryegrass, Bermudagrass, Zoysiagrass, St. Augustinegrass, Bahiagrass and Centipedegrass: Apply 3.2 to 4.0 pounds of product per 1,000 square feet of turfgrass. A second or follow-up application may be required for dense infestations of weeds or for adverse seasonal conditions. The second application of 3.2 to 4.0 pounds of product per 1,000 square feet should be used 30 days after the first application.

The maximum application rate to turf is 0.3 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year. A uniform distribution of product on the treated area is essential. Choose a calm day to ensure uniform coverage and avoid spreading granules onto Flowers, Vegetables, and Ornamental Shrubbery. Do not exceed the application rate for any area and use care in the application of this product within the drip line of Trees, Shrubs and other Ornamentals.

BROADLEAF WEEDS			
Annual Yellow Sweetclover	Clover, Sweet Clover, White Cockle	Narrowleaf Vetch Nettle	Spotted Spurge Spurweed
Aster	Cocklebur	Orange Hawkweed	Stinging Nettle
Austrian Fieldcress	Common Mullein	Oriental Cocklebur	Strawberry, India Mock
Bedstraw	Creeping Jenny	Oxalis	Tall Nettle
Beggarticks	Cudweed	Parsley-Piert	Tall Vervain
Betony, Florida	Curly Dock	Parsnip	Tansy Ragwort
Bindweed, Field	Daisy, English	Pearlwort	Tansy Mustard
Bird Vetch	Daisy, Fleabane	Pennycress	Tanweed
Bitter Wintercress	Daisy, Oxeye	Pepperweed	Thistle
Bittercress, Hairy	Dandelion	Pigweed	Trailing Crownvetch
Bitterweed	Dichondra	Pineywoods	Tumble Mustard
Black-Eyed Susan	Dogfennel	Bedstraw	Tumble Pigweed
Black Medic	Dollarweed	Plains Coreopsis	Velvetleaf
Black Mustard	False Dandelion	Plantain	Venice Mallow
Blackseed Plantain	Falseflax	Poison Ivy	Virginia Buttonweed
Blessed Thistle	False Sunflower	Poison Oak	Virginia Creeper
Bloodflower	Fiddleneck	Pokeweed	Virginia Pepperweed
Milkweed	Blue Lettuce	Poorjoe	Wavyleaf Bullthistle
	Blue Vervain	Prairie Sunflower	Western Clematis
	Bracted Plantain	Pickly Lettuce	Western Salsify
	Bristly Oxtongue	Prickly Sida	White Mustard
	Broadleaf Dock	Prostrate Knotweed	Wild Aster
	Broadleaf Plantain	Prostrate Spurge	Wild Buckwheat
	Broomweed	Prostrate Vervain	Wild Carrot
	Buckhorn	Puncturevine	Wild Four-O'Clock
	Buckhorn Plantain	Purslane,	Wild Garlic
	Bulbous Buttercup	Common	Wild Geranium
	Bull Thistle	Ragweed	Wild Lettuce
	Bullnettle	Red Sorrel	Wild Marigold
	Burclover	Hedge Bindweed	Wild Mustard
	Burdock	Hedge Mustard	Wild Onion
	Burning Nettle	Henbit	Wild Parsnip
	Burweed	Hoary Cress	Wild Radish
	Buttercup	Hoary Plantain	Wild Rape
	Buttonweed	Hoary Vervain	Wild Strawberry
	Canada Thistle	Hornsettle	Wild Sweet Potato
	Carolina Geranium	Jimsonweed	Wild Vetch
	Carpetweed	Knawel	Woodsorrel
	Catchweed	Knotweed	Wooly Croton
	Bedstraw	Kochia	Wooly
	Catnip	Lambsquarters	Morningglory
	Catsear	Lespedeza	Wooly Plantain
	Chickweed,	Mallow	Wormseed
	Common	Matchweed	Yarrow
	Chickweed,	Mexicanweed	Yellow Rocket
	Mouseear	Milk Vetch	Yellowflower
	Chicory	Morningglory	Pepperweed
	Cinquefoil	Mouseear	and other
	Clover, Crimson	Hawkweed	Broadleaf Weeds
	Clover, Hop	Mugwort	
	Clover, Red	Musk Thistle	
	Clover, Strawberry	Mustard	
		Narrowleaf Plantain	
		Spotted Catsear	

SPREADER GUIDE ONE BAG WILL COVER UP TO 10,000 SQUARE FEET

These settings can be used only as a guide, they are not absolute. The speed at which the applicator walks, spreader condition and the texture of the turf will alter the rate of application. If product remains in spreader after application to recommended square foot coverage, spread remainder over same area going at right angles (See Diagram).



SPREADER	SPREADER SETTINGS
ferti-lome® by Spyker	3.8
ferti-lome® Plus Iron Broadcast/Earthway 2850	14
ferti-lome® Plus Iron Prof. Broadcast/Earthway 2800A	14
ferti-lome® Deluxe Broadcast/Earthway 2050 S.U.	14
Earthway Drop 7300B	17
Scott Broadcast/Speedy Green	6
Republic/Scotts Broadcast	5
Republic Drop	6
Precision Broadcast/Red Devil	10
Precision Drop FH-22D	9

The above spreader settings are approximate. Calibrate your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

25-0-4 GUARANTEED ANALYSIS	
Total Nitrogen (N)	25%
2% Ammoniacal Nitrogen	
23% Urea Nitrogen	
Soluble Potash (K ₂ O)	4%
Boron (B)	0.02%
Copper (Cu)	0.05%
0.05% Water Soluble Copper	
Iron (Fe)	0.10%
0.10% Water Soluble Iron	
Manganese (Mn)	0.05%
0.05% Water Soluble Manganese	
Zinc (Zn)	0.05%
0.05% Water Soluble Zinc	
Derived from:	Sulfate of Ammonia, Urea, Muricate of Potash, Sodium Borate, Copper Sulfate, Iron Sulfate, Manganese Sulfate, Zinc Sulfate
	F370

LIMITED WARRANTY AND DISCLAIMER

The Manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use. THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the Manufacturer or the Seller. Any damages arising from a breach of the Manufacturer's Warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law. The terms of this Limited Warranty and Disclaimer cannot be varied in any written or verbal statements or agreements. No employee or agent of the Seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.



9-07

Distributed By:
VOLUNTARY PURCHASING GROUPS, INC.
P.O. BOX 460 • HWY. 56 WEST
BONHAM, TEXAS 75418
Visit Us At: www.fertilome.com

EPA Reg. No. 2217-603-7401 • EPA Est. No. 7401-TX-1

Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals>